

AD-A229 777

DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

2a. SECURITY CLASSIFICATION AUTHORITY NA		1b. RESTRICTIVE MARKINGS NA									
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA		3. DISTRIBUTION/AVAILABILITY OF REPORT  Distribution Unlimited									
4. PERFORMING ORGANIZATION REPORT NUMBER(S)  Washington University		5. MONITORING ORGANIZATION REPORT NUMBER(S)  NA									
6a. NAME OF PERFORMING ORGANIZATION  Washington University	6b. OFFICE SYMBOL (If applicable)  NA	7a. NAME OF MONITORING ORGANIZATION  Office of Naval Research									
6c. ADDRESS (City, State, and ZIP Code) Contract & Grant Administration One Brookings Drive, Campus Box 1054 St. Louis, Missouri 63130		7b. ADDRESS (City, State, and ZIP Code)  800 N. Quincy Street Arlington, Virginia 22217-5000									
8a. NAME OF FUNDING/SPONSORING ORGANIZATION  Office of Naval Research	8b. OFFICE SYMBOL (If applicable)  ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  N00014-90-J-1393									
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, Virginia 22217-5000		10. SOURCE OF FUNDING NUMBERS <table border="1"> <tr> <td>PROGRAM ELEMENT NO.</td> <td>PROJECT NO.</td> <td>TASK NO.</td> <td>WORK UNIT ACCESSION NO.</td> </tr> <tr> <td></td> <td></td> <td>4414020</td> <td></td> </tr> </table>		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.			4414020	
PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.								
		4414020									
11. TITLE (Include Security Classification)  Molecular Mechanism of Voltage-Dependent Ion Channels (U)											
12. PERSONAL AUTHOR(S)  Marshall, Garland Ross											
13a. TYPE OF REPORT  Annual	13b. TIME COVERED FROM 12/1/89 TO 11/30/90	14. DATE OF REPORT (Year, Month, Day) 11/29/90	15. PAGE COUNT 8								
16. SUPPLEMENTARY NOTATION											
17. COSATI CODES <table border="1"> <tr> <th>FIELD</th> <th>GROUP</th> <th>SUB-GROUP</th> </tr> <tr> <td></td> <td></td> <td></td> </tr> </table>		FIELD	GROUP	SUB-GROUP				18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Peptaibol, ion channel, alamethicin, emerimicin, sodium channel, alpha-helix, 3 <sub>10</sub> -helix, voltage-dependent gating. JS			
FIELD	GROUP	SUB-GROUP									
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  Please see reverse.											
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION (U)									
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Igor Vodyanoy		22b. TELEPHONE (Include Area Code) 703-696-4056	22c. OFFICE SYMBOL ONR								

DTIC  
ELECTE  
DEC 06 1990  
S E D

Item 19 - Abstract

The focus of this research is the interaction of a molecule at the lipid/water interface of a membrane with the external electric field, using peptaibols and the S4 segment of the sodium channel as model systems. Experiments are underway to assess the role of the molecular dipole in sensing the imposed molecular field; the role of  $\alpha$ -helix- $3_{10}$ -helix transitions in responding to the imposed molecular field as the transduction mechanism; and the role of the channel mouth in determining the ion selectivity associated with channel opening. Ac-(Lys(alamethicin 1-17))<sub>4</sub>-NH<sub>2</sub>, tetramethicin, which was prepared by coupling several small fragments to tetralysine while attached to a solid support, showed an effective concentration dependence 2.7 times that of alamethicin. Other oligomeric pores are in preparation using aza-crown ether scaffolds. This concentration dependence is dramatically different from the control, monomeric alamethicin-1-17 benzyl ester, which is severalfold less than alamethicin itself. This placed constraints on any model for the alamethicin channel. Analogs of emerimicin of different lengths have been prepared to probe the effect of helix length on pore stability. NMR has been used to show that a nonapeptide fragment of emerimicin changes its conformation to a predominately  $3_{10}$ -helix from the alpha-helix seen in the crystal when dissolved in DMSO. This is consistent with a theoretical analysis of the forces stabilizing the two helical forms. Umbrella sampling of molecular dynamics simulations has shown that the activation energy barrier between the  $3_{10}$ -helix and the alpha-helix is small, and that the relative stabilities of the two helical forms are a function of helix length. Solid state NMR techniques (REDOR) are being developed to allow the precise measurement of interatomic distances to determine conformations of peptaibol antibiotics in membranes. Integrations of the results from this multidisciplinary approach will provide the foundation for realistic models of voltage-gated synthetic pores.

**"Molecular Mechanism of Voltage-Dependent Ion Channels"**

**Garland R. Marshall, Ph.D.  
Principal Investigator**

**Contract N00014-90-J-1393  
R&T Code 4414020**

**Department of Pharmacology  
Washington University  
660 South Euclid Avenue  
St. Louis, Missouri 63110**

**ANNUAL REPORT  
November 29, 1990**



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## Research Objectives

### *General*

To gain an understanding of the interaction of a molecule at the lipid/water interface of a membrane with the external electric field, using peptaibols and the S4 segment of the sodium channel as a model system.

### *Specific*

To assess the role of the molecular dipole in sensing the imposed molecular field; the role of  $\alpha$ -helix-3<sub>10</sub>-helix transitions in responding to the imposed molecular field as the transduction mechanism; and the role of the channel mouth in determining the ion selectivity associated with channel opening.

### Progress (Year 1)

Significant progress has occurred in three areas, synthetic, experimental and theoretical.

### *Synthetic*

Ac-(Lys(alamethicin 1-17))<sub>4</sub>-NH<sub>2</sub>, tetramethicin, which was prepared by coupling several small fragments to tetralysine while attached to a solid support, showed an effective concentration dependence 2.7 times that of alamethicin. This concentration dependence is dramatically different from the control monomeric alamethicin 1-17 benzyl ester, which is severalfold less than alamethicin itself. Because of difficulties in determining the absolute purity of tetramethicin, we decided to prepare the same material by a totally different synthetic route which was more conducive to monitoring purity. Dr. Karol Kociolek prepared the Boc-Lys(alamethicin 1-17) monomer in solution by fragment condensation. This monomer is being used in solid phase synthesis to prepare oligomers of this peptide. In this case, any synthetic problem will differ from the desired product by 18 amino acids, which should allow easy purification by size exclusion chromatography. A second area of progress has been in the resolution of  $\alpha$ -methyl-Cys(S-benzyl) by Drs. Tom Leplawy and Urszula Slomczynska. We plan to use this compound to cross-link alamethicin sidechains. The availability of Boc-Lys(alamethicin 1-17) monomer will allow unambiguous interpretation of the position of the cross-link.

In an effort to determine the length dependence of activity, a series of emerimicin analogs has been prepared in which the emerimicin nonapeptide, Ac-Phe-MeA-MeA-MeA-Leu-Gly-Val-MeA-MeA-OBzl, has been extended by the dipeptide, Ala-MeA-OBzl, to yield the following peptides:

Ac-Phe-MeA-MeA-MeA-Leu-Gly-Val-MeA-MeA-Ala-MeA-OBzl (11)

Ac-Phe-MeA-MeA-MeA-Leu-Gly-Val-MeA-MeA-Ala-MeA-Ala-MeA-OBzl (13)

Ac-Phe-MeA-MeA-MeA-Leu-Gly-Val-MeA-MeA-Ala-MeA-Ala-MeA-Ala-MeA-OBzl (15)

These compounds will be sent for assay on bilayers when their characterization is complete.

In the past, synthetic difficulties associated with low yields in the couplings of  $\alpha,\alpha$ -dialkylamino acids have prevented the routine use of solid phase peptide synthesis in the preparation of peptaibol antibiotics. Recent developments in new coupling reagents (BOP, TBTU, etc.) have sufficiently improved the speed of reaction and yields in solution when using these sterically hindered residues that we intend to reinvestigate the use of solid phase synthesis in the preparation of analogs. Success would dramatically increase our synthetic capacity to explore structure-activity relations.

### *Experimental*

On the experimental front, we continue to explore new solid state NMR techniques (REDOR) in collaboration with their developer, Dr. Jacob Schaeffer, which we believe will be applicable in liposomes. We demonstrated this approach on an emerimicin fragment using  $^{15}\text{N}$  to perturb the  $^{13}\text{C}$  label. To enhance the sensitivity, we have decided to use  $^{19}\text{F}$  to perturb the  $^{13}\text{C}$  label and to use an adjacent  $^{15}\text{N}$  to allow selection of the label and elimination of natural abundance  $^{13}\text{C}$  signals. To this end, the following new labelled emerimicin fragments have been prepared:

FCH<sub>2</sub>CO-Phe-MeA-MeA-MeA- $^{13}\text{C}$ O- $^{15}\text{N}$ -Leu-Gly-Val-MeA-MeA-OBzl

FCH<sub>2</sub>CO-Phe-MeA-MeA-MeA-Leu-Gly(C1,C2- $^{13}\text{C}$ )-Val-MeA-MeA-OBzl

FCH<sub>2</sub>CO-Phe-MeA-MeA-MeA- $^{13}\text{C}$ O-Leu-Gly-Val-MeA-MeA-OBzl.

These peptides are being used to calibrate the REDOR approach.

Dr. Denise Beusen has completed the study of the conformation of emerimicin 1-9 benzyl ester in DMSO. Numerous technical problems had to be overcome because of the absence of  $\alpha$ -protons in much of the sequence. Dr. Beusen employed systematic search procedures developed by our group to allow assignment of hydrogen bonding partners in an unambiguous way. The structure determined is predominately  $3_{10}$ -helix, in contrast with the conformation in the crystal where this peptide shows  $\alpha$ -helical hydrogen bonding. This result complements our theoretical results, indicating a length dependence on helical preference as well as a rationale for environmental selection of helical type. A paper describing this work is nearing completion.

### *Theoretical*

On the theoretical front, we have investigated the transition between the  $3_{10}$ - and  $\alpha$ -helix by molecular dynamics. Using umbrella sampling and varying the stretch of the decapeptide, we have been able to show that the activation energy barrier between the two helical forms is relatively low compared with  $kT$ . This implies that hypotheses which consider a transition of helical forms are plausible. This is important for shorter peptides, such as emerimicin, in which the length of 15 residues is not sufficient to span the bilayer unless it assumes a  $3_{10}$ -helix. Other studies on peptides of different numbers of residues suggest that a peptide of 7 residues would show no preference between an  $\alpha$ - and  $3_{10}$ -helix conformation in a hydrophobic environment with a shorter length favoring a  $3_{10}$ -helix and a longer length favoring an  $\alpha$ -helix. While  $3_{10}$ -helices have been observed in protein crystal structures, they are all short (less than six residues in length). By providing a theoretical rationale for a facile transition between helix types, new mechanisms for other biological systems become plausible. For example, in the case of the insulin receptor, the extracellular receptor domain is connected to the tyrosine kinase domain by a single transmembrane helix. A single change in the hydrogen bonding scheme at the junction with the receptor domain on ligand binding could trigger a change in helix type and trigger a conformational change in the tyrosine kinase resulting in activation.

## Work Plan (Year 2)

In year two, we will complete the synthesis and characterization of trimethicin, tetramethicin, pentamethicin and hexamethicin using the Boc-Lys(alamethicin 1-17) monomer prepared by Dr. Kociolek. We will incorporate resolved  $\alpha$ -methyl-Cys(S-benzyl) into Boc-Lys(alamethicin 1-17) monomers and use these to prepare oligomers in which the monomers are cross-linked by disulfides as well as by the C-terminal oligoLys bridge. We will attempt to prepare cyclic oligomers of tetramethicin by synthesis on a benzyl ester support and cyclization of the oligoLys backbone between the N-terminal amine and the C-terminal carboxyl groups. If these compounds retain activity, then considerable constraints on the conformation of the active pore will have been established.

Other cyclic scaffolds will be used to build oligomers of peptaibols. We have derivatized aza-crown ethers by acylating the nitrogens with Boc- $\beta$ -alanine. Peptide fragments will be added to the amino groups of the  $\beta$ -alanine to give a tetramer in the case of the aza-crown-4 system which is our model. By increasing the size of the ring and number of nitrogen sites, one can control both channel diameter as well as the size of the mouth.

We plan to study the conformation of emerimicin itself by NMR with several solvents of varying dielectric constants to confirm the ease of transition between  $3_{10}$ - and  $\alpha$ -helix. In addition, we will initiate studies of our labelled emerimicin fragments in various solvents in the frozen state for the same reasons.

## Inventions (Year 1)

We are in the process of evaluating any commercial potential for the compounds prepared and the routes of synthesis. While no patents have been applied for, we believe that some should be in the future.

## Publications and Reports (Year 1)

Beusen, D.D. and G.R. Marshall. The generation of three-dimensional structures from NMR-derived constraints. In: *Protein Structure and Engineering*, O. Jardetzky, Ed., Plenum, pp. 97-109 (1989).

Marshall, G.R., E.E. Hodgkin, D.A. Langs, G.D. Smith, J. Zabrocki and M.T. Leplawy. Factors governing helical preference of peptides containing

multiple  $\alpha,\alpha$ -dialkyl amino acids. *Proc. Natl. Acad. Sci. USA* 87:487-491 (1990).

Leplawy, M.T., K. Kociolek, A.S. Redlinski, U. Slomczynska, J. Zabrocki, J.B. Dunbar, Jr. and G.R. Marshall. Chemical-enzymatic synthesis of emerimicin IV and III, membrane-active pentadecapeptide antibiotics. In: *Proc. 11th Am. Peptide Symp.* J. Rivier and G.R. Marshall, Eds., ESCOM, Leiden, pp. 493-495 (1990).

Marshall, G.R., D.D. Beusen, J.D. Clark, E.E. Hodgkin, J. Zabrocki and M.T. Leplawy. Helical preferences of peptides containing multiple  $\alpha,\alpha$ -dialkyl amino acids. In: *Proc. 11th Am. Peptide Symp.* J. Rivier and G.R. Marshall, Eds., ESCOM, Leiden, pp. 873-876 (1990).

Marshall, G.R., D.D. Beusen, K. Kociolek, A.S. Redlinski, M.T. Leplawy, Y. Pan and J. Schaefer. Determination of a precise interatomic distance in a helical peptide by REDOR NMR. In: *Proc. 11th Am. Peptide Symp.* J. Rivier and G.R. Marshall, Eds., ESCOM, Leiden, pp. 608-609 (1990); *J. Am. Chem. Soc.* 112:963-966 (1990).

Beusen, D.D., H. Iijima and G.R. Marshall. Structures from NMR distance constraints. *Biochem. Pharmacol.* 40:173-175 (1990).

Marshall, G.R., D.D. Beusen, J.D. Clark and E.E. Hodgkin. Helical transitions in peptides as possible gating mechanisms. Abstract, International Workshop on Transmembrane Electric Fields, Maryland, October 23-26, 1990.

Clark, J.D., D.D. Beusen, E.E. Hodgkin and G.R. Marshall. Helical transitions in peptides. Contribution for a volume to be dedicated to Prof. G.N. Ramachandran; to be published by the Indian Academy of Sciences, Bangalore, India (in press).

Hodgkin, E.E., J.D. Clark, K.R. Miller and G.R. Marshall. Conformational analysis and helical preferences of normal and  $\alpha,\alpha$ -dialkyl amino acids. *Biopolymers* (in press).

Slomczynska, U., J. Zabrocki, M.T. Leplawy and G.R. Marshall. Toward facilitated synthesis of peptaibols: Alamethicin via enzymatic segment condensation. In: *Peptides 1990*, E. Giralt and D. Andreu, Eds., ESCOM, Leiden (in press).



Leplawy, T., Jr., U. Slomczynska, M.T. Leplawy and G.R. Marshall. Peptides containing  $\alpha,\alpha$ -disubstituted amino acids. Experiments related to  $\alpha$ -methylcysteine. In: *Peptides 1990*, E. Giralt and D. Andreu, Eds., ESCOM, Leiden (in press).

Beusen, D.D., W.C. Hutton, J.J. Kotyk, J. Zabrocki, M.T. Leplawy and G.R. Marshall.  $^{13}\text{C}$  and  $^1\text{H}$  resonance assignments and structural determination of an emerimicin peptide in DMSO. In: *Peptides 1990*, E. Giralt and D. Andreu, Eds., ESCOM, Leiden (in press).

### **Training Activities**

None.

### **Awards/Fellowships**

None.

Final and Technical Reports

ADMINISTRATORS

Dr. Igor Vodyanoy, Code 1141SB (2 copies)  
Scientific Officer, Biophysics  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314

Administrative Contracting Officer  
ONR Resident Representative  
(address varies - obtain from contract or your business office)

Dr. Robert J. Nowak, Code 1113ES  
Scientific Officer, Electrochemical  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000

Program Manager  
Biological/Human Factors Division  
Code 125  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217-5000

Program Manager, Defense Technical Support  
Technology Directorate  
Office of Naval Technology, Code 223  
800 N. Quincy Street  
Arlington, VA 22217-5000

Annual and Final Reports Only (one copy each)

DoD ACTIVITIES

Commander  
Chemical and Biological Sciences Division  
Research Army Research Office, P. O. Box 1221  
Research Triangle Park, NC 27709

Head  
Biomolecular Engineering Branch  
Code 6190  
Naval Research Laboratory  
Washington, DC 20375

Directorate of Life Sciences  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, DC 20332